

100-104 / 4-18 / MDDC-984

MDDC-984

MDDC - 984

UNITED STATES ATOMIC ENERGY COMMISSION

184-INCH CYCLOTRON VERTICAL BEAM OSCILLATIONS
IN THE REGION OF 82-INCH RADIUS

by
James Vale


University of California

This document consists of 7 + 1 pages

Date of Manuscript: February 26, 1947


Date Declassified: May 23, 1947

Its issuance does not constitute authority
to declassify copies or versions of the
same or similar content and title
and by the same author(s).



Technical Information Division, Oak Ridge Directed Operations
AEC, Oak Ridge, Tenn., 10-22-48--1000-11980

Printed in U.S.A.
PRICE 10 CENTS



Digitized by the Internet Archive
in 2011 with funding from
University of Florida, George A. Smathers Libraries with support from LYRASIS and the Sloan Foundation

184-INCH CYCLOTRON VERTICAL BEAM OSCILLATIONS IN THE REGION OF 82-INCH RADIUS

By James Vale

Experiment done by: L. Houser, R. Watt, and J. Vale

INTRODUCTION

It was discovered early in the operation of the 184-inch cyclotron, that the beam did not go out to the expected 85-inch radius.

The Theoretical Group suggested that the beam did not go beyond a radius of 82 inches because of vertical oscillations building up in the vicinity of this point. According to their calculations, the vertical oscillations built up a large enough amplitude at the point where $n = 0.2$ ($n = -R/H \, dH/dR$) so that the beam hit the dee and was lost. An attempt had been made to find this vertical spreading, previously, but it was unsuccessful. Consequently, the following experiment was performed.

EXPERIMENTAL SETUP

Several targets were made of 1/16-inch copper sheet. These were U-shaped and were placed in the tank and bombarded. The plane of the sheet was vertical, and the arms of the U were roughly parallel to the dee face. See sketch of target appended hereto.

Thus, the beam had to traverse the slot in the U to reach the back of the target. Hence, any vertical spreading in the beam would result in its hitting the U arms.

These targets were run into the tank, one at a time, on the regular probe mechanism, were bombarded with a large deuteron beam for about 1 to 3 minutes, removed from the tank and radioautographs were taken to determine where the beam was hitting. The targets differed only in the width of the slot. These slot widths varied from 2 1/2 to 4 1/2 inches.

RESULTS

The autographs indicate a rapid spreading vertically of the beam at about 81 1/2 inches. This agrees quite closely with the point at which $n = 0.2$ from magnetic measurements. The autographs indicate that the vertical spreading is large enough so that most of the beam gets caught by the dee at this point. (Prints of autographs appear on following pages.)




Figure 1. Sketch of target.




Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.

UNIVERSITY OF FLORIDA



3 1262 08909 7231